

OFFICE INFORMATION

Procedure date:

Arrival time:

Location:

Bowel Preparation for Colonoscopy With MoviPrep

Colonoscopy is a routine, generally safe, pain-free procedure used to examine the colon. There are steps that you, the patient, must take before the procedure to ensure the highest level of safety and effectiveness.

- 1 Make arrangements to have someone drive you home after the procedure
 - Colonoscopy involves sedation, and you will not be allowed to leave unaccompanied
- 2 Follow your physician's instructions regarding medicines to avoid and diet to follow before your procedure
- 3 Finish the entire bowel prep regimen as described on the following page

MoviPrep is available at most retail pharmacy outlets. If your pharmacy does not have MoviPrep in stock, it can be ordered and delivered within 24 hours.

Information for your pharmacist: NDC 65649-201-75



MoviPrep®
(PEG-3350, Sodium Sulfate, Sodium Chloride, Potassium Chloride,
Sodium Ascorbate and Ascorbic Acid for Oral Solution)

**Low-Volume Prep.
High-Volume Efficacy.**

Patient Instructions


The MoviPrep carton contains 4 pouches and a disposable container for mixing. You must complete the entire prep to ensure the most effective cleansing.

Beginning at:

STEP 1


- Empty 1 Pouch A and 1 Pouch B into the disposable container
- Add lukewarm drinking water to the top line of the container. Mix to dissolve

If preferred, mix solution ahead of time and refrigerate prior to drinking. The reconstituted solution should be used within 24 hours.



STEP 2

- The MoviPrep container is divided by 4 marks. Every 15 minutes, drink the solution down to the next mark (approximately 8 oz), until the full liter is complete
- Drink 16 oz of the clear liquid of your choice



Beginning at:

- Repeat steps 1 and 2

If you have any questions, please call our office at _____.

MoviPrep is indicated for cleansing of the colon as a preparation for colonoscopy in adults 18 years of age or older. MoviPrep is contraindicated in patients who have had a severe hypersensitivity reaction to any of its components. MoviPrep should be used with caution in patients using concomitant medications that increase the risk of electrolyte abnormalities, in patients with known or suspected hyponatremia, severe ulcerative colitis, ileus, gastrointestinal obstruction or perforation, gastric retention, toxic colitis, toxic megacolon, or glucose-6-phosphate dehydrogenase deficiency. In clinical trials, abdominal distension, anal discomfort, thirst, nausea, and abdominal pain were the most common adverse reactions to MoviPrep administration.

MoviPrep contains a maximum of 2.33 mg of phenylalanine per treatment.

Please see accompanying full Prescribing Information for MoviPrep.



Web site: www.salix.com 1700 Perimeter Park Drive, Morrisville, NC 27560 Tel • 866.669.SLXP (7597)
©2006 Salix Pharmaceuticals, Inc. All rights reserved. Printed in USA. MOV 06/04

MoviPrep[®]
(PEG-3350, Sodium Sulfate, Sodium Chloride, Potassium Chloride,
Sodium Ascorbate and Ascorbic Acid for Oral Solution)

Low-Volume Prep.
High-Volume Efficacy.